


Ciprofloxacin Suspension for Babies

How to give Ciprofloxacin Suspension To Your Child


- Mix the medicine as shown to the right.
- Shake the medicine for about 15 seconds before measuring out the next dose.
- After it is mixed, keep the medicine in the refrigerator. It may be kept covered in the refrigerator for 14 days.
- **Give two doses each day. Give one in the morning and one in the evening.**
- Measure the amount of liquid medicine for your child's dose, as marked below.
- To make it taste better, the suspension can be mixed with a small amount of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**

Child's weight	Give this much	How often each day
7-12 pounds (3-5 kg)	1 ml	Two times
13-22 pounds (6-10 kg)	2 ml	Two times
23-33 pounds (11-15 kg)	3 ml	Two times
34-44 pounds (16-20 kg)	4 ml	Two times


Mixing Instruction:




1. The small bottle contains the medicine. **Do not chew the medicine.** Mix the medicine with the liquid in the larger bottle.



2. Open both bottles. Press down on the cap as you twist it to the left.



3. Pour the medicine into the larger bottle of liquid. **Do not add water or any other liquid.** Be sure you get all the medicine out of the small bottle.




4. Close the large bottle. Be sure the cap is on tight. Shake the bottle well for about 15 seconds. The medicine is ready for use.

Sida Hooraha Ciprofloxacin Suspension Loo Siiyo Ilmahaaga


- Dawada u qas sida lagu muujiyay midigta.
- Rux dawada qiyaas ah 15 sekan (il-biriqsi) ka hor cabirista qadarka qaadashada ee ku xiga.
- Ka dib marka la qaso, dawada ku hay qaboojiyaha. Iyada oo daboolan waxa lagu hayn karaa qaboojiyaha mudo ah 14 maalmood.
- **Sii laba qadar qaadasho maalin kasta. Hal sii subixii halka kalena fiidkii.**
- Sida hoos lagu calaamadeeyay, u cabir qadarka qaadashada ilmahaaga ee dawada hooraha ah.
- Si dhadhankeeda loo fiicneeyo, waxa hooraha lagu qasi karaa qadar yar oo ah cunto ama miir khudaar. **Hubso in ilmuhu qaato dhammaan isku-darka si uu u helo qadar buuxa oo ah dawada.**

Miisaanka ilmaha	Sii qadarkan	Intee jeer maalin kasta
3-5 kg (7-12 rodol)	1 ml	Laba wakhti
6-10 kg (13-22 rodol)	2 ml	Laba wakhti
11-15 kg (23-33 rodol)	3 ml	Laba wakhti
16-20 kg (34-44 rodol)	4 ml	Laba wakhti


Farriimaha Qasidda




1. Dhalada yar waxa ku jira dawada. **Ha calaanjin dawada.** Dawada ku qas hooraha ku jira dhalada weyn.



2. Fur labada dhalo. Hoos u riix furka adiga oo u wareejinaya bidix.



3. Dawada ku shub dhalada weyn ee hooruhu ku jiro. **Ha ku darin biyo ama hoore kale.** Hubso inaad dhammaan dawada ku shubtid dhalada yar.



4. Xidh dhalada weyn. Hubso in furku ku adag yahay. Si fiican u rux dhalada qiyaas ah 15 sekan (il-biriqsi). Dawadu waa u diyaar in la isticmaalo.