



# Giving Doxycycline Pills to Your Child



- The amount of medicine depends on the child’s weight.
- For a child who weighs less than 55 pounds (25kg), you should use the Doxycycline Suspension for your child, not the pills. Please refer to that page for directions.
- **Give two doses each day. Give one in the morning and one in the evening.** Give the amount marked below for your child.
- To make it taste better, the medicine can be crushed and mixed with a **small amount** of food or juice. **Be sure the child takes all this mixture to get the full dose of medicine.**

Child’s weight	Morning Dose Amount	Evening Dose Amount
Under 55 pounds (under 25 kg)	See the page about Doxycycline Suspension for medicine directions	
55-80 pounds (25-36 kg)	½ pill	½ pill
81-98 pounds (37-44 kg)	1 pill	½ pill
99 pounds or more (45kg or more)	1 pill	1 pill



## 儿童服用Doxycycline药片

- 药物剂量依儿童体重而定。
- 体重不足25公斤（55磅），应使用Doxycycline悬浮液，勿使用药片。请参阅有关说明。
- **每天服用两次。早晚各一次。**按下表的药量给孩子服药。
- 为了改善药物的味道，可将药片弄碎后与**少量**食物或果汁混合后服用。**务必**要让孩子将混合物全部服下，以确保全剂量服药。

儿童体重	早晨剂量	晚上剂量
25公斤以下（55磅以下）	参阅Doxycycline悬浮液服用说明	
25-36公斤（55-80磅）	½片	½片
37-44公斤（81-98磅）	1片	½片
45公斤或以上（99磅或以上）	1片	1片