

Nitroglycerin

Nitroglycerin is a medicine used to treat chest pain called angina. **Take nitroglycerin as directed by your doctor.**

- If you are pregnant or breastfeeding, talk to your doctor before using this medicine.
- Always keep your nitroglycerin with you.
- This medicine may make you dizzy or lightheaded when you stand up or get out of bed. Get up slowly from a sitting or lying position.
- Ask your doctor if it is safe for you to drink alcohol while using nitroglycerin medicine.
- Keep nitroglycerin in its original bottle to keep its strength. The medicine should cause a sting or a tingle under your tongue. If not, you may need a new bottle of medicine.
- Keep the medicine out of the reach of children.

If you use tablets:

1. Sit down and rest 2 to 3 minutes.
2. If you have chest pain or discomfort that is not better with rest, place 1 tablet under your tongue. Do not swallow the tablet.
3. If you still have chest pain after 5 minutes, put a second tablet under your tongue.
4. If the chest pain does not go away after another 5 minutes, use a third tablet. You may take up to 3 tablets in 15 minutes.
5. If you still have pain, **call 911**. Do not drive yourself to the hospital because you may be having a heart attack.

If you use spray:

1. Sit down and rest for 2 to 3 minutes.
2. If you have chest pain or discomfort that is not better with rest, hold the can up with the spray button on top as close to your mouth as you can.

3. Press down on the button and spray the medicine onto your tongue or under it. Don't swallow right away. Wait 10 seconds.
4. If your chest pain lasts longer than 5 minutes, spray a second dose into your mouth.
5. If the chest pain does not go away after another 5 minutes, spray a third dose. You may use up to 3 sprays in 15 minutes.
6. If you still have pain, **call 911**. Do not drive yourself to the hospital because you may be having a heart attack.

Call your doctor if you have any of these medicine side effects:

- Flushed or hot face
- Nausea or vomiting

Call your doctor or 911 right away if you have any of these side effects:

- Trouble breathing
- Increased chest pain
- Dizziness, fainting or severe weakness
- Severe headaches
- Itching or swelling
- Blurred vision
- Dry mouth

Talk to your doctor, nurse or pharmacist if you have any questions or concerns about taking nitroglycerin.

2005 – 9/2010 Health Information Translations

Unless otherwise stated, user may print or download information from www.healthinfotranslations.org for personal, non-commercial use only. The medical information found on this website should not be used in place of a consultation with your doctor or other health care provider. You should always seek the advice of your doctor or other qualified health care provider before you start or stop any treatment or with any questions you may have about a medical condition. The Ohio State University Medical Center, Mount Carmel Health System, OhioHealth and Nationwide Children's Hospital are not responsible for injuries or damages you may incur as a result of your stopping medical treatment or your failure to obtain medical treatment.